

CE/CSE 5710: Modeling & Simulation

The Inventory Problem

February 2, 2012

Consider the inventory problem. Inventory is stocked so that a store does not fall below demand at any given time. Problem definition is as follows:

- Replenishment of stock takes place at the end of each time period labeled $n = 0, 1, 2, \dots$
- Total average demand during any period n is the stationary random variable ϵ_n . Hence,

$$Pr\{\epsilon_n = k\} = a_k \quad \text{for } k = 0, 1, 2, \dots \quad (1)$$

$$\text{where } a_k \geq 0 \quad \text{and} \quad \sum_{k=0}^{\infty} a_k = 1 \quad (2)$$

- A replenishment policy is established as (s, S) where $S > s$ - i.e., the stock is replenished at the end of each time period n iff it dips below s to the level S . The stock is checked at the end of each time period.
- X_n denotes the stock size at the end of each time period n just before the stock is replenished (or not). $\{X_n\}$ is considered to be a stochastic process defined on the space $(-\infty, S]$.

Therefore the following relationships hold:

$$X_{n+1} = \begin{cases} X_n - \epsilon_{n+1}; & \text{if } s < X_n \leq S \\ S - \epsilon_{n+1}; & \text{if } X_n \leq s \end{cases}$$

If the successive demands $\epsilon_1, \epsilon_2, \dots$ are independent random variables then the stock values X_0, X_1, X_2, \dots constitute a Markov chain whose transition probability matrix can be stated to be:

$$P_{i,j} = Pr\{X_{n+1} = j | X_n = i\} = \begin{cases} Pr\{\epsilon_{n+1} = i - j\}; & \text{if } s < i \leq S \\ Pr\{\epsilon_{n+1} = S - j\}; & \text{if } i \leq s \end{cases}$$

Problem

Using the above formulation, consider a spare parts inventory model in which either 0, 1 or 2 spare parts are demanded in a period with $Pr\{\epsilon_n = 0\} = 0.5$, $Pr\{\epsilon_n = 1\} = 0.4$, $Pr\{\epsilon_n = 2\} = 0.1$.