

CE3501 Environmental Engineering
Part II. Environmental Biology
Fall 2005

Homework Assignment #2, Due Monday, 10/24 in class

1. Balance the following redox reactions:


- a. Sulfate reduction: $\text{CH}_2\text{O} + \text{SO}_4^{2-} \leftrightarrow \text{HCO}_3^- + \text{HS}^- + \text{H}^+$
- b. Nitrification: $\text{NH}_4^+ + \text{O}_2 \leftrightarrow \text{NO}_3^- + \text{H}_2\text{O} + \text{H}^+$
- c. Iron reduction: $\text{CH}_2\text{O} + \text{Fe}(\text{OH})_{3(s)} \leftrightarrow \text{HCO}_3^- + \text{Fe}^{2+} + \text{OH}^- + \text{H}_2\text{O}$
- d. Aerobic oxidation of MTBE: $\text{C}_5\text{H}_{12}\text{O} + \text{O}_2 \leftrightarrow \text{CO}_2 + \text{H}_2\text{O}$

You may use either the method of half-reactions or balance the whole reaction equation.

2. Write and balance the half-reactions (oxidation, reduction) for the following reactions:

- a. Manganese reduction: $\text{CH}_2\text{O} + \text{MnO}_2 + \text{H}^+ \leftrightarrow \text{HCO}_3^- + \text{Mn}^{2+} + \text{H}_2\text{O}$
- b. Denitrification: $\text{NO}_3^- + \text{CH}_2\text{O} \rightarrow \text{N}_2 + \text{HCO}_3^- + \text{H}^+ + \text{H}_2\text{O}$

3. What is the theoretical oxygen demand (ThOD) for complete oxidation of a saturated solution of the pesticide malathion? The solubility of malathion is 144 mg/L. (**223 mg/L**)



4. What is the average oxidation state of carbon in malathion ($\text{C}_{10}\text{H}_{19}\text{O}_6\text{PS}_2$)? The oxidation state of the sulfur is -2 and that of phosphorus is +5. (**Ans: +0.8**)

5. A tank truck of milk runs off of a bridge and dumps its entire contents (16 m^3) into a river. If the k_L for degradation of the milk is 0.4 d^{-1} , how long would be required to reduce the BOD in the river by 99% of the value at the spill site? Assume no tributaries or other water sources dilute the milk in the river as it flows downstream from the spill site. (**ANS: 11.5 d**)

6. How much oxygen is *actually* consumed during decomposition of a waste depends on the extent to which the waste is biodegradable, i.e. amenable to oxidation by microbes. Ammonia is totally biodegradable and thus the nitrogenous ThOD (theoretical) and the nitrogenous BOD (actual) are equal. Not so for carbonaceous compounds; the carbonaceous BOD (actual) may be significantly less than the carbonaceous ThOD (theoretical) if the compound is poorly degradable. The 5-day BOD test provides an estimate of the amount of oxygen that a waste will actually consume. The malathion waste in Problem 3 was found to have a 5-day BOD of 125 mg/L and a rate constant (k_L) of 0.2 d^{-1} . Calculate the ultimate BOD of the solution. Is the malathion very biodegradable? What fraction of the potential oxidation actually occurs? (**198 mgO₂/L, Yes, 88%**).